

01/783896

WEST**Freeform Search**

Database: **US Patents Full-Text Database** ▲
 US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
 IBM Technical Disclosure Bulletins ▼

Term: 14 and two step\$1 ▲

Display: 10 Documents in Display Format: CIT Starting with Number 1

Generate: ☐ Hit List ☒ Hit Count ☐ Side by Side ☐ Image

Search

Clear

Help

Logout

Interrupt

Main Menu

Show 8 Numbers

Edit 8 Numbers

Preferences

Cases

Search HistoryDATE: Thursday, February 14, 2002 [Printable Copy](#) [Create Case](#)Set Name Query
side by sideHit Count Set Name
result set

DB=USPT,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ

L6 14 and two step\$14 L6L5 L453 L5L4 immuno same polymerase chain reaction\$153 L4L3 immuno same polymerase chain reaction same two step\$10 L3L2 immuno same polymerase chain reaction same two same step\$1 same
RNA same cDNA0 L2L1 immuno near5 polymerase chain reaction near5 two step\$10 L1

END OF SEARCH HISTORY

A rectangular button with a thin black border and the text "Generate Collection" centered inside.

L6: Entry 2 of 4

File: USPT

Feb 23, 1999

DOCUMENT-IDENTIFIER: US 5874226 A

TITLE: In situ immunodetection of antigens

Brief Summary Paragraph Right (5):

Another known method of immunodetection takes advantage of the immuno-PCR (Polymerase Chain Reaction) methodology (2). The PCR method is similar to the Cantor method up to the incubation with biotinylated DNA, however, instead of using multiple rounds of streptavidin and biotinylated DNA incubation, the DNA/biotin/streptavidin/antibody complex is washed out with a low pH or high salt buffer that releases the antibody. The resulting wash solution is then used to carry out a PCR reaction with suitable primers with appropriate controls. At least in theory, the enormous amplification capability and specificity of PCR can be utilized to detect a single antigen molecule.

Detailed Description Paragraph Right (11):

Other amplification technologies are also applicable, including two steps, 3 steps, PAP, and APAAP methods.